

# Effectiveness of 2% chlorhexidine gel compared to two solvents commonly used in endodontic retreatment

Claudiani Saraiva **CAETANO**<sup>1</sup>

Máira do **PRADO**<sup>2</sup>

Brenda Paula Figueiredo de Almeida **GOMES**<sup>3</sup>

Ezilmara Leonor Rolim de **SOUSA**<sup>4</sup>

## ABSTRACT

**Objective:** The present study has compared the action of 2% chlorhexidine gel to two solvents commonly used in endodontic retreatment, xylol and eucalyptol, regarding the cleaning of the root canal walls. **Methods:** Forty-five human single-rooted teeth were randomly divided into three groups. The teeth were instrumented, filled and radiographed before being stored in an oven at 37°C for 60 days. The filling material was removed from root canals according to the following techniques: Kerr and Hedström files with xylol (G1), eucalyptol (G2), and 2% chlorhexidine gel (G3). After removing the filling material, the teeth were radiographed in ortho- and mesio-radial directions. The radiographs were analyzed by three double-blinded examiners for the presence of remaining

filling material. Next, the teeth were longitudinally fractured and the resulting faces were digitally scanned. The Image Tool software was used to assess the amount of remaining filling material, and data were statistically analyzed. **Results:** With regard to the radiographic analysis, no statistically significant differences were observed between the groups studied (i.e. chlorhexidine, xylol and eucalyptol). However, when digitalized images were analyzed, xylol was found to be significantly more efficient for cleaning the root canals compared to 2% chlorhexidine gel. **Conclusion:** It can be stated that xylol was the most effective solution for removal of filing material compared to 2% chlorhexidine gel and eucalyptol.

**Keywords:** Retreatment. Chlorhexidine. Solvents. Root canal preparation.

**How to cite this article:** Caetano CS, Prado M, Gomes BPFA, Sousa ELR. Effectiveness of 2% chlorhexidine gel compared to two solvents commonly used in endodontic retreatment. *Dental Press Endod.* 2012 Oct-Dec;2(4):15-9.

» The authors report no commercial, proprietary or financial interest in the products or companies described in this article.

<sup>1</sup>Graduation Student, School of Dentistry, UFPel.

<sup>2</sup>Associate Professor, UFRJ

<sup>3</sup>Professor, School of Dentistry of Piracicaba, UNICAMP.

<sup>4</sup>Adjunct Professor, School of Dentistry, UFPel.

Received: November 19, 2012. Accepted: November 25, 2012.

Contact address: Máira do Prado  
Av. Limeira 901 – Vila Areão  
CEP: 13.414-903 – Piracicaba/SP – Brazil  
Email: mairapr@hotmail.com

## Introduction

Removal of filling material and cleaning of root canals are considered important procedures for a successful endodontic retreatment.<sup>1,2</sup> In order to remove the filling material from the root canals, different techniques have been proposed by the literature. These techniques include the use of rotary or manual instruments in association or not with solvents.<sup>3</sup>

Different solvents have been proposed for aiding in the removal of filling material from root canals, among them one can cite chloroform, eucalyptol and xylol.<sup>3,4</sup> Although effective, most endodontists do not use solvents because of the toxic action of such substances on periapical tissues.<sup>5,6</sup>

According to Oliveira,<sup>7</sup> chlorhexidine gel can be an alternative to the use of solvents. In fact, this substance has low toxicity<sup>8</sup> and allows the root canal walls to be mechanically cleaned due to its viscosity, thus compensating the action of solvents.<sup>8</sup> In addition, chlorhexidine has a wide-spectrum antibacterial activity.<sup>9,10,11</sup> According to Gomes et al,<sup>10</sup> 2% chlorhexidine gel has shown to be highly effective against *Enterococcus faecalis*, which is a relevant fact as Sundqvist et al,<sup>12</sup> Molander et al<sup>13</sup> and Pinheiro et al<sup>14</sup> reported that this microorganism is associated with cases of endodontic failure.

The objective of the present study was to compare the action of 2% chlorhexidine gel to two solvents largely used in endodontic retreatment, xylol and eucalyptol, regarding the cleaning of the root canal walls.

## Material and Methods

The Human Research Ethics Committee of the Federal University of Pelotas, Dentistry School (process number 012/2006) has approved this study. A total of 45 human single-rooted teeth (incisors and canines) with complete apices were used for study, all being stored in saline solution (Basa – Indústria Farmacêutica Basa Ltda, Caxias do Sul, RS, Brazil) and kept at constant temperature of 37°C in an oven. By using a digital calliper, the roots had their length standardised to 15 mm. Double-faced diamond discs (KG Sorensen, São Paulo, SP, Brazil) mounted on a micro-motor and straight handpiece (Kavo Extra –Torque 605C, Brazil) were used for this standardization.

The foramens were standardised by using a Flexofile #15 file (Dentsply Maillefer, Petrópolis, RJ, Brazil) and the working length was set at 1 mm short of the apex.

The root canals were instrumented according to the step-back technique, which consisted in preparing the coronal third with Gates-Glidden burs #2 and #3 (Dentsply Maillefer, Petrópolis, RJ, Brazil). For apical preparation, the apical stop was standardised to calliper #35 and step back performed in 1 mm increments until #55 file. During instrumentation, the root canals were irrigated with 2.5% sodium hypochlorite solution (Vida Nova Farmácias de Manipulação). After instrumentation, EDTA (Iodontosul Indústria Odontológica do Sul) was used for 3 minutes for removal of the smear layer and then a final irrigation with saline solution was applied.

Root canals were dried with paper tips (Endpoints) and filled according to the lateral condensation technique by using gutta-percha cones and Endofill sealer (Dentsply). The teeth were radiographed in the mesial-distal and buccal-lingual orientations for analysis of the quality of the obturation.

Next, the root canal entry was temporarily restored with intermediate restorative material (Dentsply) and the teeth were stored in an oven at 37°C during 60 days to allow the sealer to set.

After 60 days, the restorative material was removed, including a coronal 5 mm of filling material (gutta-percha + sealer) by using Gates-Glidden burs #2 and #3 in order create a reservoir for the auxiliary chemical agent being used.

The teeth were divided into 3 groups (n = 15) depending on the auxiliary chemical agent used during the preparation:

- Group 1: Manual instrumentation with #15 to #45 K-files (Dentsply Maillefer) followed by #15 to #45 Hedström files (Dentsply Maillefer) in association with the use of 0.5 mL of 2% chlorhexidine gel at each instrumentation and abundant irrigation with 3 ml of saline solution (Basa).
- Group 2: Manual instrumentation with #15 to #45 K-files (Dentsply Maillefer, ) in association with #15 to #45 Hedström files (Dentsply Maillefer), adding 0.5 mL of xylol (Merck at each instrumentation and final irrigation with 3 ml of saline solution (Basa).
- Group 3: Manual instrumentation with #15 to #45 K-files (Dentsply Maillefer) in association with #15 to #45 Hedström files (Dentsply Maillefer), adding 0.5 mL of eucalyptol (Biodinâmica) at each instrumentation and final irrigation with 3 ml of saline solution (Basa).

The maximum time for removing the filling material from each root canal was defined in 30 minutes. Removal of the filling material was considered complete when remnants of gutta-percha or filling material recovering the instruments were no longer observed, with these criteria already being proposed by Hulsmann, Stotz<sup>15</sup> and Imura et al.<sup>16</sup>

After removing the filling material, the teeth were radiographed (Spectro 70X-Dabi-Atlante) at a focal distance of 200 mm with an X-ray unit operating at 70 KVp, 8mA, and exposure time of 0.5 seconds. The radiographs taken were in the mesial-distal and buccal-lingual directions.

The radiographs were evaluated by three double-blinded examiners for the presence of remnants of gutta-percha, attributing scores described by Ferreira et al.<sup>17</sup> (Table 1) for the different thirds (coronal, middle and apical) regarding the amount of radiopaque debris.

After the radiographic analysis, the teeth were longitudinally sectioned in the buccal-lingual direction by using double-faced diamond discs (KG Sorensen) and surgery chisel (Neumar). Image of the segments were taken in pairs (semi-parts) by digitalizing the image with a scanner device HP Deskjet F300 All-In-One Series (HP-Brazil) operating at a 1200 dpi resolution and presence of a stainless steel ruler for image standardization. Next, the software Image Tool was used to analyze the images and helped calculate the sum of all areas (regions) where remnants of the filling material were present. For this analysis, the measurements were in mm<sup>2</sup>.

Kruskal-Wallis' test was used ( $p < 0.05$ ) for statistical analysis of the amount of radiopaque debris (radiographic analysis of the scores), whereas ANOVA ( $p < 0.05$ ) and Tukey's tests ( $p < 0.05$ ) were used to analyze the sum of the areas containing remnants of filling material.

## Results

With regard to the radiographic analysis of the scores, no statistically significant difference was observed between the three groups studied, that is, chlorhexidine, xylol, and eucalyptol (Table 2). However, when the total area of remaining filling material was assessed with the Image Tools software, one could observe a statistically significant difference between the chlorhexidine and xylol groups, with the latter being more efficient than 2% chlorhexidine gel for cleaning the root canals. No statistically significant difference was found regarding the use of eucalyptol and 2% chlorhexidine gel.

With regard to the analysis of the thirds, Figure 1 shows the scores obtained from each of them. It was observed a higher degree of cleaning efficiency in the coronal third, followed by middle and apical thirds.

## Discussion

According to Wilcox and Swift,<sup>18</sup> a successful endodontic retreatment is strongly associated with the cleaning of the root canal walls. The present study has evaluated the root canal cleaning with different substances by means of radiographic analysis and digital imaging.




Figure 1. Scores attributed to the different thirds.

Table 1. System of scores attributed to radiographic analysis according to the amount of radiographic debris.

Scores	Significant
0	Absence of radiopaque debris
1	Less than 25% of debris
2	Between 25% and 50% of debris
3	More than 50% of debris

Table 2. Values of the sum of remaining filling material present in the root canals in the three groups.

Groups	Radiographic analysis (scores)	Image analysis (Area of remaining material)
Chlorhexidine	0.8 ± 1.1 <sup>a</sup>	6.1 ± 4.2 <sup>b</sup>
Xilol	0.3 ± 0.5 <sup>a</sup>	2.5 ± 3.0 <sup>a</sup>
Eucalyptol	0.8 ± 0.9 <sup>a</sup>	5.4 ± 4.9 <sup>b</sup>

Notes: Letters a and b mean statistically significant differences.

In the present work, the root length was standardised to 15 mm with the removal of the coronal portion of the teeth. This procedure was performed according to Al-Omari and Dummer,<sup>19</sup> who state that such a removal allows the root canal to be better viewed morphologically and possible interferences eliminated during the retreatment procedures.

Although several previous studies had shown that the time required for removing the filling material is around 20 minutes, regardless of the technique used,<sup>2,16</sup> we have opted to define a time of 30 minutes.

With regard to the different analyses used in the present study, the radiographic one was used because it is a methodology already established in the literature. According to Tanomaru Filho et al,<sup>20</sup> the conventional radiographic technique using periapical films in two radiographic orientations provides good conditions for analysis. However, according to Ferreira et al,<sup>17</sup> it is not possible to assess the amount of remaining filling material existing on the root canal walls by using periapical films.

In the present study, according to such a technique, it was observed a better cleaning of the coronal third compared to the middle and apical ones. This finding may be associated to the use of Gates-Glidden burs for this third, enabling better cleaning compared to the middle and apical thirds as these were cleaned with manual instruments only.<sup>7</sup> Despite also being described by Wilcox<sup>21</sup> and Ferreira et al,<sup>17</sup> these burs were used here to prepare a reservoir for storing the chemical substances used.

With regard to the analysis of the root canal cleaning using digital imaging, this technique was shown to be efficient for quantifying the amount of remaining filling material, which was also reported by Bramante and Betti.<sup>22</sup>

In the present work, when this technique was employed it was observed a better cleaning of the dentinal walls in association with the use of xylol compared to 2% chlorhexidine gel and eucalyptol. In addition, in the case of the former substance, it was difficult to reach the entire working length in some of the teeth, as higher pressure was applied to the files in order to remove the filling material from the root canals. This happened because chlorhexidine does not act directly on the filling material, that is, it does not alter the properties of this material by causing its

dissolution. In fact, because of its viscosity, this substance only allows the filling material to be removed in fragments, which are dislocated from the inside of the root canals. Also, due to the need to exert higher pressure to the instrument, the use of manual files in association to 2% chlorhexidine gel might cause a deviation of the root canal, which did not occur in the present study as the teeth being used already had wide and straight root canals.

The findings reported in the present study are in accordance with Oyama et al,<sup>23</sup> who evaluated the properties of several solvents by assessing the weight loss of gutta-percha cones following the action of the substances at different times, concluding that xylol was the most effective. Additionally, Bueno and Valdrighi<sup>24</sup> reported better results with the use of xylol compared to eucalyptol. However, our findings differ from those found by Oliveira,<sup>7</sup> who reported that chlorhexidine was better than the other solvents studied. This discrepancy may be associated to the type of instrumentation, since Oliveira<sup>7</sup> had used rotary instruments that soften the gutta-percha by heating it, whereas in the present study manual files were used instead.

## Conclusion

According to the methodology used and the results found in the present work, one can conclude that the use of xylol has favoured the removal of filling material as well as the cleaning of dentinal walls compared to the use of eucalyptol and 2% chlorhexidine gel.

## References

1. Aun CE, Santos M. Quantity of apical extruded material and efficiency of five different methods of removing gutta-percha and sealer from root canals: "in vitro" evaluation. *Rev Fac Odontol FZL*. 1989;1(2):63-73.
2. Friedman S, Moshnov J, Trope M. Efficacy of removing glass ionomer cement, zinc oxide eugenol, and epoxy resin sealers from retreated root canals. *Oral Surg Oral Med Oral Pathol*. 1992;73(5):609-12.
3. Lopes HP, Siqueira Jr JF. *Endodontia, biologia e técnica*. Rio de Janeiro: Guanabara Koogan; 2004.
4. De Deus Q. *Endodontia*. 5ª ed. Rio de Janeiro: Medsi; 1992.
5. Barbosa SV, Burkard DH, Spångberg LS. Cytotoxic effects of gutta-percha solvents. *J Endod*. 1994;20:6-8.
6. Vajrabhaya LO, Suwannawong SK, Kamolroongwarakul R, Pewklieng L. Cytotoxicity evaluation of gutta-percha solvents: Chloroform and GP-Solvent (limonene). *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2004;98(6):756-9.
7. Oliveira DP. Avaliação "in vitro" da remoção de material obturador empregando diferentes técnicas durante o retratamento endodôntico [dissertação mestrado]. Piracicaba: Faculdade de Odontologia da Unicamp; 2002.
8. Lee TH, Hu CC, Lee SS, Chou MY, Chang YC. Cytotoxicity of chlorhexidine on human osteoblastic cells is related to intracellular glutathione levels. *Int Endod J*. 2010;43:430-5.
9. Ferraz CC, Gomes BP, Zaia AA, Teixeira FB, Souza-Filho FJ. In vitro assessment of the antimicrobial action and the mechanical ability of chlorhexidine gel as an endodontic irrigant. *J Endod*. 2001;27:452-5.
10. Gomes BPFA, Ferraz CCR, Vianna ME, Beber VB, Teixeira FB, Souza-Filho FJ. In vitro antimicrobial activity of several concentrations of sodium hypochlorite and chlorhexidine gluconate in the elimination of *Enterococcus faecalis*. *Int Endod J*. 2001;34(6):424-8.
11. Sena NT, Gomes BP, Vianna ME, Berber VB, Zaia AA, Ferraz CC, et al. In vitro antimicrobial activity of sodium hypochlorite and chlorhexidine against selected single-species biofilms. *Int Endod J*. 2006;39(11):878-85.
12. Sundqvist G, Figdor D, Persson S, Sjögren U. Microbiologic analysis of teeth with failed endodontic treatment and the outcome of conservative retreatment. *Oral Surg Oral Med Oral Pathol*. 1998;85(1):86-93.
13. Molander A, Reit C, Dahlen G, Kvist T. Microbiological status of root-filled teeth with apical periodontitis. *Int Endod J*. 1998;31(1):1-7.
14. Pinheiro ET, Gomes BPFA, Ferraz CCR, Sousa ELR, Teixeira FB, Souza-Filho FJ. Microorganisms from canals of root filled teeth with periapical lesions. *Int Endod J*. 2003;36(1):1-11.
15. Hulsmann M, Stolz S. Efficacy, cleaning ability and safety of different devices for gutta-percha removal in root canal retreatment. *Int Endod J*. 1997;30(4):227-33.
16. Imura N, Kato AS, Hata GI, Uemura M, Toda T, Weine F. A comparison of the relative efficacies of four hand and rotatory instrumentation techniques during endodontic retreatment. *Int Endod J*. 2000;33(4):361-6.
17. Ferreira JJ, Rhodes JS, Pitt Ford TR. The efficacy of gutta-percha removal using profiles. *Int Endod J*. 2001;34(4):267-74.
18. Wilcox L, Swift M. Endodontic retreatment in small and large curved canals. *J Endod*. 1991;17(7):313-5.
19. Al-Omari MA, Dummer PMH. Canal blockage and debris extrusion with eight preparation techniques. *J Endod*. 1995;21(3):154-8.
20. Tanomaru Filho, Leonardo MR, Silva LAB, Castro ET. Avaliação radiográfica in vitro da capacidade de limpeza de técnicas de retratamento endodôntico. *Rev Assoc Paul Cir Dent*. 1999;53(3):238-41.
21. Wilcox L.R. Endodontic retreatment with halotane versus chloroform solvent. *J Endod*. 1995;21(6):305-7.
22. Bramante CM, Betti LV. Efficacy of Quantec rotatory instruments for gutta-percha removal. *Int Endod J*. 2000;33(5):463-7.
23. Oyama KON, Siqueira EL, Santos M. Ação de diferentes solventes sobre os cones de guta percha. *ECLER Endod*. 1999;1(3):1-8.
24. Bueno CES, Valdrighi L. Efetividade de solventes e de técnicas na desobturação dos canais radiculares. Estudo in vitro. *Rev ABO Nac*. 2000;8(1):21-5.