

Suggesting a new therapeutic protocol for traumatized permanent teeth: Case report

Jefferson J. C. **MARION**¹

Laiza Vicente **MARTELOSSO**²

Juliana Yuri **NAGATA**³

Thiago Farias R. **LIMA**⁴

Adriana de Jesus **SOARES**⁵

ABSTRACT

Introduction: The case reported herein consists of two teeth of one single patient, initially immobilized with a semi-rigid retainer used for 15 days without endodontic intervention. **Objective:** The aim of this article is to report a case of two teeth with extrusive luxation treated with a filling paste of calcium hydroxide, 2% chlorhexidine gel and zinc oxide. **Methods:** After a one-month follow-up, patient's teeth did not present evidence of pulp vitality. Radiographic examination revealed signs of external resorption. Endodontic treatment was carried out in association with a new treatment protocol using intracanal

dressing applied in one single session and remaining in the root canal for four years. **Results:** The filling paste remained in the root canal for 24 months without being replaced. The case presented improvements in periapical lesions without inflammatory resorption. **Conclusion:** The filling paste proves to be successful and effective in treating traumatized teeth with root resorption.

Keywords: Endodontics. Calcium hydroxide. Tooth socket. Extrusive luxation. Dental trauma. Chlorhexidine gel. Zinc oxide.

How to cite this article: Marion JJC, Martellosso LV, Nagata JY, Lima TFR, Soares AJ. Suggesting a new therapeutic protocol for traumatized permanent teeth: Case report. *Dental Press Endod.* 2014 Jan-Apr;4(1):71-7. doi: <http://dx.doi.org/10.14436/2178-3713.V4.N1.071-077.oar>.

¹PhD in Clinical Dentistry/Endodontics, State University of Campinas (UNICAMP). Professor, Department of Endodontics, Brazilian Dental Association (ABO), Maringá. Assistant professor, Ingá College.

²Degree in Dentistry, Ingá College.

³PhD in Clinical Dentistry/Endodontics, UNICAMP. Assistant professor, Department of Endodontics, Federal University of Sergipe (UFS).

⁴PhD in Clinical Dentistry/Endodontics, State University of Campinas (UNICAMP).

⁵PhD in Clinical Dentistry/Endodontics, State University of Campinas (UNICAMP). Visiting professor, Dental Trauma Service, UNICAMP.

» Patients displayed in this article previously approved the use of their facial and intraoral photographs.

» The authors report no commercial, proprietary or financial interest in the products or companies described in this article.

Submitted: November 11, 2013. Revised and accepted: November 02, 2013.

Contact address: Jefferson J. C. Marion
Rua Néo Alves Martins, 3176 – 6º andar – Sala 64 – Centro
CEP: 87-013.060 – Maringá/PR – Brazil — E-mail: jefferson@jmarion.com.br

Introduction

There is a high incidence of dental trauma in the general population. It may occur in adults and children, but it is more common among children and young adults. Dental trauma is mainly caused by fall, collision, bicycle or car accidents and sports practice, all of which may lead to a wide variety of dental trauma.¹⁻⁴ A total of 85.39% of incisor trauma occurs probably due to their positioning in the arch.^{5,6}

Traumatic injury is classified under a variety of names, among which luxation is the most common. It includes trauma of tooth displaced from its original position, in which case it is classified according to the direction in which the tooth has been displaced, namely: concussion, subluxation, extrusive luxation, lateral luxation, intrusive luxation and avulsion.^{7,8} Damages caused by trauma not only affect hard tissues and dental pulp, but also affect supporting tissues alone or in combination. In extreme cases, trauma may lead to tooth loss.⁶ Extrusive luxation is the partial displacement of a tooth, causing it to come out of the socket. According to Andreassen et al,^{9,10} it accounts for 64% of dental trauma cases. Complications arising from this type of trauma may occur weeks or years after the accident.¹¹

Calcification, pulp necrosis and root resorption are the most common sequelae of trauma.^{7,12} The latter has the worst prognosis^{13,14} and, according to the American Association of Endodontics,¹⁵ it is associated with physiological and pathological processes that result in dentin, cementum and alveolar bone loss. Additionally, it may be classified as inflammatory or replacement resorption.¹⁶

Several authors¹⁷⁻²⁰ assert that extrusive luxation may be treated by repositioning extruded incisors and using a semi-rigid immobilization, both of which do not interfere in occlusion or function. Immobilization must be kept for approximately 15 days, according to the individual characteristics of each case. Retention is performed with orthodontic wire and light-cured resin and it aims at regenerating periodontal fibers. Also, it is extremely important that the clinician perform patient's clinical as well as radiographic follow-up, as cases of pulp necrosis and inflammatory root resorption require endodontic therapy.

Endodontic treatment of reimplanted teeth require the use of intracanal dressing to supplement disinfection and restrain or ease the inflammatory process of

resorption. The procedure most commonly used before root canal filling is the use of calcium hydroxide periodically changed and associated with different vehicles. This intracanal dressing is used for having excellent antimicrobial activity and inhibiting the action of cells involved in root resorption.²¹⁻³¹

A new therapy associating calcium hydroxide, 2% chlorhexidine gel and zinc oxide has been recently proposed for treatment of avulsed teeth.^{32,33,34} The combination of these substances results in a provisional filling paste that remains in the root canal for a long period of time, thereby eliminating the need for restorative procedures and yielding satisfactory results,^{32,35,36} except for cases of replacement resorption which are a continuous process.³² This filling paste provides the patient with comfort, requires less visits to the dentist, and proves less expensive as it does not need to be replaced. In cases of incomplete root formation, it promotes apical closure and relieves clinical signs and symptoms of traumatized teeth for a period of 9 months.³³ Furthermore, it promotes periapical repair in teeth with inflammatory root resorption.³³ Such properties are explained by its high capacity of diffusing throughout the root dentin, inhibiting bacterial growth in the outer root surfaces^{37,38} and, therefore, yielding satisfactory results.

Thus, the aim of this article is to report a case of two teeth with extrusive luxation treated with a filling paste of calcium hydroxide, 2% chlorhexidine gel and zinc oxide.

A case report

A 17-year-old male patient sought the Service of Dental Trauma at the School of Dentistry, Piracicaba (FOP-UNICAMP) with dental history of trauma in teeth #11 and #21. Emergency assistance was provided at the Santa Casa hospital of Limeira/SP where extrusive luxation of teeth #11 and #21 was diagnosed. Dental trauma occurred as a result of sports practice. A semi-rigid retainer was installed and kept for 15 days. Subsequently, the patient was referred to FOP-UNICAMP for further treatment.

He sought the services of FOP-UNICAMP a month after the trauma had occurred. His initial clinical and radiographic exams revealed teeth #11 and #21 with no pulp vitality, pain at vertical/horizontal percussion or palpation. Periapical radiograph (Fig 1) revealed complete root formation, external root resorption and

periapical lesion of traumatized teeth. Since it was a case of pulp necrosis, endodontic treatment was recommended and carried out with a putty paste made of calcium hydroxide associated with 2% chlorhexidine gel and zinc oxide (2:1:2).

After crown opening and complete isolation, the septic-toxic medication was neutralized and biomechanical preparation of the crown-apex was carried out with Gates Gliden bur #5,4,3 (Dentsply/Maillefer, Ballaigues, Switzerland) with a view to decontaminating the cervical and middle third of teeth #11 and #21. Odontometry was performed with an electronic apical locator (Novapex, Forum Technologies, RishonLeZion, Israel). Root canals underwent manual instrumentation to their working length with #45 files (Dentsply/Maillefer, Ballaigues, Switzerland). During the instrumentation procedure, 2% chlorhexidine gel (Endogel, Essencial Pharma, Itapetininga/MG, Brazil) was inserted into the root canal at each change of instrument, followed by irrigation with 5 ml of saline solution.

Smear layer was removed by irrigation with 3 ml of 17% EDTA for 3 minutes, followed by final irrigation with saline solution. Root canals were dried with absorbent paper points (Konne Indústria e Comércio de Materiais Odontológicos Ltda., Belo Horizonte/MG, Brazil) and filled with a paste of calcium hydroxide P.A. (Biodinâmica Quim. e Farm. Ltda., Iporã/PR, Brazil) associated with 2% chlorhexidine gel (Endogel, Essencial Pharma, Itapetininga/MG, Brazil) and zinc oxide (S.S. White Artigos Dentários, Ltda., Rio de Janeiro/RJ, Brazil). The paste was coltosol or putty-consistent and was prepared in a 2:1:2 ratio. It was inserted by increments with medium and fine medium vertical condenser (Konne Indústria e Comércio de Materiais Odontológicos Ltda., Brazil) throughout the entire root canal. Afterwards, a periapical radiograph was taken to ensure that the filling procedure had been properly performed. Root canals were then sealed with coltosol (Vigodent S/A Indústria e Comércio, Rio de Janeiro/RJ, Brazil) and composite resin (Filtek Z350, 3M Dental Products, Saint Paul, USA) (Fig 2).

Every 3 months, the patient went back to the university for clinical and radiographic follow-up sessions that revealed the presence of intracanal medication/paste completely filling the canal, thus eliminating the need for removal. After 12 months of treatment, the clinician observed remission of periapical lesion (Fig 3A).

After 24 months of follow-up, exams revealed the intracanal medication filling the entire root canal, thus eliminating the need for removal. Periapical lesion a remission and interruption of inflammatory resorption were observed (Fig 3B).

Discussion

Endodontic treatment is a predictable way of preserving a tooth in cases of dental trauma. Treatment success and high-quality immediate restoration aim at restoring patient's esthetics and function for years. Thus, endodontic treatment is considered a safe and feasible option.³⁹

Cases in need of endodontic treatment due to trauma have led to different therapeutic protocols used to minimize potential sequelae. The use of sodium hydroxide,⁴⁰ polymyxin B-Otosporin,⁴¹ Lysozymes,⁴² formocresol,⁴³ chlorhexidine^{44,45,46} and calcium hydroxide, in different associations^{47,48}, has been reported.



Figure 1. Initial radiograph 30 days after trauma and after removal of semi-rigid retainer.




Figure 2. A) Calcium hydroxide P.A.; 2% chlorhexidine gel (Endogel) and zinc oxide used as intracanal dressing in a 2:1:2 ratio. B) Final consistency. C) Final radiograph revealing the quality of insertion of the intracanal dressing. D) Vertical condenser used to insert the intracanal dressing.




Figure 3. A) 12-month follow-up radiograph. B) 24-month follow-up radiograph.

Some authors^{24,49,50} recommend that calcium hydroxide be used as intracanal dressing applied with periodic changes and at different time intervals, given that it is considered the most effective intracanal medication of all.^{51,52,53} According to Pacios et al,⁵⁴ calcium hydroxide yields satisfactory clinical results and may be used in association with different vehicles, for instance, distilled water, chlorhexidine, propylene glycol, anaesthetic solution, PMCC and PMCC + propylene glycol. However, this case report did not implement periodic changes and, for this reason, disagrees with the aforementioned studies. Conversely, it corroborates Felipe et al³⁵ who asserts that there is no need for applying calcium hydroxide past with periodic changes in teeth with incomplete root formation and incomplete root canal systems.

Even though calcium hydroxide is the most used intracanal medication, it cannot be considered as universal, given that it is not effective against all types of bacteria present in root canal systems.⁵⁵ Since chlorhexidine presents highly satisfactory antimicrobial properties, it is important that it be used in association with calcium hydroxide. Calcium hydroxide associated with chlorhexidine aims at enhancing the antimicrobial properties of the former, keeping its biological characteristics as well as its mechanisms of physical barrier.⁴⁷ Zinc oxide is a yellowish, white, thin, odorless, amorphous, insoluble in water or ethanol, radiopaque and slightly antiseptic powder⁵⁶ that may be present in endodontic cement⁵⁷ and gutta-percha cones.⁵⁸

Based on the aforementioned benefits, this case report aimed at assessing the results yielded by a therapeutic protocol using the three elements together: calcium hydroxide, 2% chlorhexidine gel and zinc oxide mixed together to form a coltosol-consistent paste that did not dissolve and could be applied in one single session, thus eliminating the need for periodic changes during trimestrial control. The outcomes of this research corroborate those of previous studies.^{32,33,59,60,61}

Due to the fact that the intracanal paste used during the procedure did not dissolve, as revealed by radiographic exams, it is suggested that complete root canal filling was guaranteed by zinc oxide, since calcium hydroxide and chlorhexidine gel had already been dissolved.^{32,59} Similar results were yielded by our study in which a 24-month follow-up was performed. During this period, the intracanal dressing did not have to be periodically changed; periapical repair was achieved and root resorption was stabilized, thereby proving the

effectiveness of the medication in the long run and eliminating the need for replacement inside the root canal. Gomes et al³¹ and Souza-Filho et al⁶² also conducted studies in which the intracanal dressing was not periodically replaced. The authors observed that during the follow-up phase, zinc oxide possibly functions as an inert material that provides root canal sealing and, as a consequence, prevents contamination and allows periapical repair. In addition to that, they concluded that the medication can remain inside the root canal for a period not greater than 4 years. Those results disagree with the studies by Moorer and Genet⁶³ who suggested that additional researches be conducted to further investigate the biological properties of zinc oxide, given that these authors do not consider it as an inert material.

The association of calcium hydroxide, 2% chlorhexidine gel and zinc oxide (2:1:2 ratio) was previously studied.^{37,38,47,62,64,65} These *in vitro* studies demonstrated the antimicrobial action of the mixture, as well as its capacity of keeping an alkaline pH and its proper consistency when inserted into the root canal. Furthermore, the association of calcium hydroxide, 2% chlorhexidine gel and zinc oxide proves to have a high diffusion capacity in the root dentin, thus causing inhibition of bacterial growth in outer root surfaces — as previously mentioned — which may have favored root resorption control.

Conclusion

Based on the results of this study and the findings of the literature, it is reasonable to conclude that:

- » The filling paste of calcium hydroxide, 2% chlorhexidine gel and zinc oxide (2:1:2 ratio) proves effective in treating traumatized permanent teeth with complete root formation and extrusive luxation. This intracanal dressing stimulates periapical lesion repair and interruption of inflammatory resorption.
- » The technique allows treatment to be performed within one single session, as the intracanal medication remains active afterwards. Additionally, it proves advantageous for the patient due to being inexpensive and having shorter chair time.
- » Due to having a cortisol-consistency, the provisional filling paste of calcium hydroxide, 2% chlorhexidine gel and zinc oxide (2:1:2 ratio) is easily inserted into the root canal. Additionally, it favors radiographic visualization.

References

1. Caldas Jr AF, Burgos ME. A retrospective study of traumatic dental injuries in a Brazilian dental trauma clinic. *Dent Traumatol*. 2001;17(6):250-3.
2. Paiva JG, Antoniazzi JH. Endodontia: bases para a prática clínica. 2a ed. São Paulo: Artes Médicas; 1991. 886 p.
3. Souza-Filho FJ, Soares AJ, Gomes BPFA, Zaia AA, Ferraz CCR, Almeida J. Avaliação das injúrias dentárias observadas no Centro de Trauma Dental da Fop-Unicamp. *RFO UPF: Rev Facul Odontol Univ Passo Fundo*. 2009;14(2):116.
4. David J, Astrom AN, Wang NJ. Factors associated with traumatic dental injuries among 12-year-old schoolchildren in South India. *Dent Traumatol*. 2009;25(5):500-5.
5. Bezerra AG, Abrão CV, Belmonte FM, Caldeira CL. Levantamento epidemiológico dos casos de traumatismos dentais atendidos no CADE-Trauma durante o ano de 2004. 13a Reunião Anual de Pesquisa da FOUFSP. RPG. São Paulo; 2004.
7. Andreasen JO, Andreasen FM. Texto e atlas colorido de traumatismo dental. 3a ed. São Paulo: Artmed; 2001.
8. Lopes HP, Siqueira JF. Endodontia biologia e técnica. 3a ed. Rio de Janeiro: Guanabara Koogan; 2010.
9. Andreasen JO. Etiology and pathogenesis of traumatic dental injuries. A clinical study: of 1298 cases. *Scand J Dent Res*. 1970;78(4):329-42.
10. Andreasen FM, Vestergaard PB. Prognosis of luxated permanent teeth: the development of pulp necrosis. *Dent Traumatol*. 1985;1(6):207-20.
11. Robertson A, Andreasen FM, Bergenholtz G, Andreasen JO, Norén JG. Incidence of pulp necrosis subsequent to pulp canal obliteration from trauma of permanent incisors. *J Endod*. 1996;22(10):557-60.
12. Hecova H, Tzigkounakis V, Merglova V, Netolicky J. A retrospective study of 889 injured permanent teeth. *Dent Traumatol*. 2010;26(6):466-75.
13. Soares AJ, Gomes BPFA, Zaia AA, Ferraz CCR, Souza-Filho FJ. Relationship between clinical radiographic evaluation and outcome of teeth replantation. *Dental Traumatol*. 2008;24(2):183-8.
14. Guan Y, Qin M. A retrospective study of pulp healing after luxation injuries. *Zhonghua Kou Qiang Yi Xue Za Zhi*. 2008;43(9):520-3.
15. Lima TFR. Análise das sequelas clínicas e radiográficas de dentes traumatizados: estudo retrospectivo [dissertação]. Piracicaba (SP): Universidade Estadual de Campinas; 2012.
16. Dumsha TC. Luxation injuries. *Dent Clin North Am*. 1995;39(1):79-91.
17. Araújo MAM, Valera MC. Tratamento clínico dos traumatismos dentários. São Paulo: Artes Médicas; 1999.
18. Andreasen JO, Andreasen FM, Bakland LK, Flores MT. Traumatic dental injuries: a manual. 2nd ed. Munksgaard; 2003.
19. Flores MT, Andersson L, Andreasen JO, Bakland LK, Malmgren B, Barnett F, et al. Guidelines for the management traumatic dental injuries II. Avulsion of permanent teeth. *Dent Traumatol*. 2007;23:130-6.
20. Di Angelis AJ, Andreasen JO, Ebeleseder KA, Kenny DJ, Trope M, Sigurdsson A, et al. International Association of Dental Traumatology. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: fractures and luxations of permanent teeth. *Dent Traumatol*. 2012;28(1):2-12.
21. Souza V, Bernabé PFE, Holland R, Nery MJ, Mello W, Otoboni Son JA. Nonsurgical treatment of teeth with periapical lesions. *Rev Bras Odontol*. 1989;46:39-46.
22. Panzarini SR, Souza V, Holland R, E. Dezan Jr. Treatment of teeth with chronic periapical lesions. Influence of different types of root canal dressing and root canal filling material. *Rev Odontol UNESP*. 1998;27(2):509-26.
23. Katebzadeh N, Hupp J, Trope M. Histological periapical repair after obturation of infected root canals in dogs. *J Endod*. 1999;25(5):364-8.
24. Trope M. Clinical management of the avulsed tooth: present strategies and future directions. *Dent Traumatol*. 2002;18(1):1-11.
25. Holland R, Otoboni Son JA, Souza V, Nery MJ, Bernabé PFE, Dezan Jr E. Periapical repair formulations with different Ca (OH)2. Study in dogs. *Rev Assoc Paul Cir Dent*. 1999;53:327-31.
26. Leonardo MR, Silveira FF, Silva LA, Tanomaru Filho M, Utrilla LS. Calcium hydroxide root canal dressing: histopathological evaluation of periapical repair at different time periods. *Braz Dent J*. 2002;13(1):17-22.
27. Tanomaru Filho M, Leonardo MR, Silva LAB. Effect of irrigating solution and calcium hydroxide root canal dressing on the repair of apical and periapical tissues of teeth with periapical lesion. *J Endod*. 2002;28(4):295-9.
28. Holland R, Souza V, Tagliavini RL, Milanezi LA. Healing process of teeth with open apices. Histological study. *Bull Tokyo Dent Coll*. 1971;12(4):333-8.
29. Bystrom A, Haaponen RP, Sundqvist G. The antibacterial effect of camphorated paramono chlorophenol, camphorated phenol and calcium hydroxide in the treatment of infected root canals. *Endod Dent Traumatol*. 1985;1(5):170-5.
30. Trope M, Moshonov J, Nissan R, Buxt P, Yesilsoy C. Short vs. long-term calcium hydroxide treatment of established inflammatory root resorption in replanted dog teeth. *Endod Dent Traumatol*. 1995;11(3):124-28.
31. Gomes BPFA, Montagner F, Berber VB, Zaia AA, Ferraz CCR, Almeida JFA, Souza-Filho FJ. Antimicrobial action of intracanal medicaments on the external root surface. *J Dent*. 2009;37(1):76-81.
32. Soares AJ. Análise clínica e radiográfica de dentes traumatizados submetidos a um protocolo de medicação intracanal com a associação hidróxido de cálcio, clorexidina gel 2% e óxido de zinco, sem trocas periódicas [tese]. Piracicaba (SP): Universidade Estadual de Campinas; 2007.
33. Soares AJ, Souza-Filho FJ. Traumatized teeth submitted to a new intracanal medication pro-tocol. *Braz J Dent Traumatol*. 2011;2(2):1-5.
34. Buck CLBP. Avaliação clínica e radiográfica de dentes reimplantados submetidos ao tratamento endodôntico utilizando a associação de hidróxido de cálcio, clorexidina gel 2% e óxido de zinco como pasta obturadora do canal em sessão única [dissertação]. Campinas (SP): Faculdade São Leopoldo Mandic; 2011.
35. Felipe MCS, Felipe WT, Marques MM, Antoniazzi JH. The effect of the renewal of calcium hydroxide paste on the apexification and periapical healing of teeth with incomplete root formation. *Int Endod J*. 2005;38(7):436-42.
36. Steiner JC, Dow PR, Cathey GM. Inducing root end closure of nonvital permanent teeth. *J Dent Child*. 1968;35(1):47-54.
37. Montagner F, Gomes BPFA, Berber VB, Zaia AA, Souza-Filho FJ. Ação antimicrobiana de medicações intracanaís na superfície radicular externa frente a diferentes microorganismos. In: Anais da 23ª Reunião da Sociedade Brasileira de Pesquisa Odontológica, 2006. Atibaia: SBPQO; 2006. p. 126. [Resumo IC 052].
38. Gomes BPFA, Montagner F, Berber VB, Zaia AA, Ferraz CCR, Almeida JFA, Souza-Filho FJ. Antimicrobial action of intracanal medicaments on the external root surface. *J Dent*. 2009;37(1):37-81.
39. Hargreaves KM. Treatment planning: comparing the restored endodontic tooth and the dental implant. *Endodontics: colleagues for excellence*. 2007 [Acesso 22 set. 2011]. Disponível em: <http://www.aae.org/dentalpro/clinicaltopics>.
40. Niwa M, Milne KC, Ribi E, Rudbach JA. Alteration of physical chemical and biological properties of endotoxin by treatment with mild alkali. *J Bacteriol*. 1969;97(3):1069-77.

41. Oliveira LD, Leão MVP, Carvalho CAT, Camargo CHR, Valera MC, Jorge AOC. In vitro effects of calcium hydroxide and polymyxin B on endotoxins in root canals. *J Dent.* 2005;33(2):104-14.
42. Ohno N, Morrison DC. Interaction of lysozyme with bacterial lipopolysaccharide. *The FASEB J: Official Publication of the Federation of American Societies for Experimental Biology.* 1988;2:680.
43. Sant'Anna AT, Ramalho LTO, Spolidorio DMP. Effect of the formocresol on bacterial LPS in mouse's subcutaneous tissue. *J Dent Res.* 2000;79:1084.
44. Silva LA, Leonardo MR, Assed S, Tonomaru FM. Histological study of the effect of some irrigation solutions on bacterial endotoxin in dogs. *Braz Dent J.* 2004;15(2):109-14.
45. Gomes IC, Chevitaes O, Almeida NS, Salles MR, Gomes GC. Diffusion of calcium through dentin. *J Endod.* 1996;22(11):590-5.
46. Tanomaru JMG, Leonardo MR, Tanomaru Filho M, Boneti Filho I, Silva LAB. Effect of different irrigation solutions and calcium hydroxide on bacterial LPS. *Int Endod J.* 2003;36(11):733-9.
47. Gomes BPFA, Vianna ME, Senna NT, Zaia AA, Ferraz CCR, Souza-Filho FJ. In vitro evaluation of the antimicrobial activity of calcium hydroxide combined with chlorhexidine gel used as intracanal medicament. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;102(4):544-50.
48. Gomes BPFA, Sato E, Ferraz CC, Teixeira FB, Zaia AA, Souza-Filho FJ. Evaluation of time required for recontamination of coronally sealed canals medicated with calcium hydroxide and chlorhexidine. *Int Endod J.* 2003; 36(4):604-9.
49. Andersson L, Andreasen JO, Day P, Heithersay G, Trope M, Diangelis AJ, Kenny DJ, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth. *Dent Traumatol.* 2012;28:88-96.
50. Leonardo MR, Silva LAB, Tonomaru Filho M, Bonifácio KC, Ito IY. In vitro evaluation of anti-microbial activity of sealers and pastes used in Endodontics. *J Endod.* 2000;26(7):391-4.
51. Heithersay GS. Calcium hydroxide in the treatment of pulpless teeth with associated pathology. *J B Endod Soc.* 1975;8(2):74-93.
52. Cvek M, Hollender L, Nord CE. Treatment of non-vital permanent incisors with calcium hydroxide. VI. A clinical, microbiological and radiological evaluation of treatment in one sitting of teeth with mature or immature root. *Odont Revy.* 1976;27(2):93-108.
53. Martin D, Crabb H. Calcium hydroxide in root canal therapy: a review. *Br Dent J.* 1977;142(9):277-83.
54. Pacios MG, de la Casa ML, de Bulacio MI, López ME. Influence of different vehicles on the pH of calcium hydroxide pastes. *J Oral Sci.* 2004;46(2):107-11.
55. Gomes BPFA, Ferraz CCR, Garrido FD, Rosalen PL, Teixeira FB, Souza-Filho FJ. Microbial susceptibility to calcium hydroxide pastes and their vehicles. *J Endod.* 2002;28(11):758-61.
56. Siqueira Jr JF, Rôças IN. Polymerase chain reaction-based analysis of microorganisms associated with failed endodontic treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;97(1):85-94.
57. Pizzo G, Giammanco GM, Cumbo E, Nicolosi G, Gallina G. In vitro antibacterial activity of endodontic sealers. *J Dent.* 2006;34(1):35-40.
58. Spangberg LSW. Instruments, materials and devices. In: Cohen S, Burns RC, editors. *Pathways of the pulp.* 7th ed. St Louis: CV Mosby; 1998. p. 508-10.
59. Soares AJ, Nagata JY, Casarin RCV, Almeida JFA, Gomes BPFA, Zaia AA, et al. Apexification by using a new intracanal medicament: a multidisciplinary case report. *Iran Endod J.* 2012;7(3):165-70.
60. Soares AJ, Lima TFR, Lins FF, Herrera Morante DR, Gomes BPFA, Souza-Filho FJ. Un nuevo protocolo de medicación intraconducto para dientes con necrosis pulpar y rizogénesis incompleta. *Rev Estomatol Hered.* 2011;21(3):145-9.
61. Buck CLBP, Soares AJ, Buck A, Nagata JY, Zaia AA, Souza-Filho FJ. Avaliação de dentes reimplantados submetidos a um novo protocolo terapêutico. *Rev Assoc Paul Cir Dent.* 2012;66(3):200-5.
62. Souza-Filho FJ, Soares AJ, Vianna ME, Zaia AA, Ferraz CC, Gomes BP. Antimicrobial effect and pH of chlorhexidine gel and calcium hydroxide alone and associated with other materials. *Braz Dent J.* 2008;19(1):28-33.
63. Moorer WR, Genet JM. Evidence for antibacterial activity of endodontic gutta-percha cones. *Oral Surg Oral Med Oral Pathol.* 1982;53(5):503-7.
64. Almeida GC, Montagner F, Berber VB, Zaia AA, Souza-Filho FJ, Gomes BPFA. Antibacterial activity of zinc-oxide-calcium hydroxide intracanal medicaments against selected endodontic pathogen [abstract 111]. *Braz J Oral Sci.* 2006;5(18):11-38.
65. Montagner F. Avaliação in vitro da ação antimicrobiana de diferentes medicações intracanal na superfície radicular externa [monografia]. Piracicaba (SP): Universidade Estadual de Campinas; 2007.